



Product Data Sheet

Cat # RP-1476

Deslorelin

Size: 5 mg

Source: Deslorelin is a potent LHRH/GnRH agonist has a molecular formula of $C_{64}H_{83}N_{17}O_{12}$, Pyr-His-Trp-Ser-Tyr-D-Trp-Leu-Arg-Pro-NHEt having a Mw of 1282.47 Dalton. Deslorelin is being studied in the treatment of cancer as a way to block sex hormones made by the ovaries or testicles. It belongs to the family of drugs called gonadotropin-releasing hormone analogs. It is used for the induction of ovulation in mares. Deslorelin binds to and activates pituitary gonadotropin releasing hormone (GnRH) receptors. Continuous, prolonged administration of goserelin in males results in pituitary GnRH receptor desensitization and inhibition of pituitary secretion of follicle stimulating hormone (FSH) and luteinizing hormone (LH), leading to a significant decline in testosterone production; in females, prolonged administration results in a decrease in estradiol production. The protein (1 mg/ml) was lyophilized with no additives.

Usage:

This item is for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

RP-1476

050412P

Applications and Suggested Dilutions: It is recommended to reconstitute the lyophilized Deslorelin in sterile 18M Ω -cm H₂O not less than 100 μ g/ml, which can then be further diluted to other aqueous solutions. Users must optimize the appropriate concentration and conditions for each assay. Greater than 99.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE. Users must optimize the appropriate concentration and conditions for each assay.

Storage and Stability: Lyophilized Deslorelin although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Deslorelin should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). **Please prevent freeze-thaw cycles.** If supplied in powder then reconstitute it in 100 μ l water for 1 mg/ml stock and store in liquid at 4°C for ~1 week or aliquots in suitable size and store at -20°C for long term storage.