ALPHA DIAGNOSTIC INTERNATIONAL

Product Specification Sheet

ATP Binding Cassette subfamily G, member 8 (ABCG8) Antibodies

Cat. # ABCG81-P	Rat ABCG8 Control/blocking peptide # 1	**SIZE**: 100 ug
Cat. # ABCG81-A	**Rabbit** Anti-Rat ABCG8 IgG # 1 (aff pure)	**SIZE**: 100 ug

The ATP binding cassette (ABC) superfamily of membrane transporters is one of the largest protein classes known, and counts for numerous proteins involved in trafficking of biological molecules across membranes, host-defense mechanism to xenobiotics. The first known members were P-glycoprotein (P-gp) and multidrug resistant protein (MRP), cause multidrug resistance when transfected into drug-sensitive cells. In addition, increasing numbers of ABC proteins have recently been identified. The human ABCG1 (ABC, subfamily G, member 1) gene encodes a member of ABC superfamily that mediates the ATP-dependent translocation of variety of amphiphilic and lipophilic molecules. ABCG proteins were able to transport substances across cellular membranes and against concentration gradient they require an input of energy, which requires the hydrolysis of ATP, directly or indirectly.

Rat ABCG8 protein is alternatively spliced: A full length isoforms 3 (694-aa), isoforms 2 (671-aa; missing 56-77 aa, and 398-aa); isoforms 1 (672aa, missing 56-77 aa). Actual expression of various protein isoforms and their physiological relevance is not understood. ABCG8 may form heterodimers with ABCG5 or be tightly coupled to ABCG5 along a pathway regulating dietary-sterol absorption and excretion. ABCG8 protein is highly expressed in liver, intestine and colon. Defects in ABCG8 cause sitosterolemia.

Source of Antigen and Antibodies

Antigen	15-aa peptide from Rat **ABCG8** (protein accession #P58248 refs 1); **Designation (ABCG81-P, control or blocking peptide**) conjugated to KLH. Epitope location ~ within C-terminus, Cytoplasmic
Ab Host/type	Rabbit, Polyclonal, Aff pure IgG (**cat # ABCG81-A**) purified over antigen-agarose column
2-ab	**Goat Anti-rabbit IgG-HRP** cat # 20320 (AP, biotin, FITC conjugates also available)
-ve control IgG	Cat # 20009-1, Control Rabbit (non-immune) Serum IgG for use as –ve control in ELISA, Western or IHC

Form & Storage of Antibodies/Peptide Control

Affinity pure IgG

100 ug/100ul solution lyophilized powder

Supplied in **Buffer**: PBS+0.1% BSA

Reconstitute powder in PBS at 1mg/ml

Control/blocking peptide

100 ug/100 ul solution lyophilized powder

Supplied in Buffer: PBS pH 7.5,

Reconstitute powder in PBS at 1 mg/ml.

Storage

Short-term: unopened, undiluted liquid vials at -20OC and powder at 4oC or -20oC..

Long-term: at –20C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at –20oC or below.
Shipping: 4oC for solutions and room temp for powder

Recommended Usage

Western Blotting (1-10 ug/ml for affinity pure antibody using ECL technique).

ELISA: Control peptide can be used to coat ELISA plates at 1 ug/ml and detected with antibodies (0.5-1 ug/ml for affinity pure).

Histochemistry & Immunofluorescence: Not tested. We recommend the use of aff pure IgG at 2-20 ug/ml.

Specificity & Cross-reactivity

Rat ABCG81-P control peptide is 100% conserved in rat ABCG-8 isoforms 1-3. It is 65% conserved in mouse ABCG8 and no appreciable conservation in human ABCG8. No significant conservation of ABCG81-P is observed with other ABCGs (ABCG1-7) or other proteins. Antibody reactivity and actual expression of various protein isoforms and their physiological relevance is not understood. Antibody cross-reactivity in various species has not been studied. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity (see detailed protocol at: the web site).

General References: Lu. K et al (2001) Am. J. Hum. Genet. 69 (2), 278-290; Berge, K. E et al (2000) Science 290 (5497), 1771-1775; Lee, M. H et al (2001) Nat. Genet. 27 (1), 79-83; L. Austin Doyle (1998) PNAS Vol. 95, 15665-15670.

*This product is for In vitro research use only.

Related material available from ADI

Antibodies for ABCG1 to ABCG8.

ABCG81-A-P 71213S

Alpha Diagnostic Intl Inc., 6203 Woodlake Center Dr, San Antonio, TX 78244, USA; **Email**: service@4adi.com
☎ (800) 786-5777; (210) 561-9515; **Fax**: (210) 561-9544; **Web Site**: www.4adi.com