

Product Specification Sheet

**Human Glut-1 Antibodies**

Cat. # GT14-P	<b>Human</b> Glut-1 control peptide # 1	<b>SIZE:</b> 100 ug
Cat. # GT14-A	<b>Rabbit</b> Anti-Human Glut-1 IgG # 1 (aff pure)	<b>SIZE:</b> 100 ug

Most mammalian cells transport glucose through a family of membrane proteins known as glucose transporters. Molecular cloning of these glucose transporters has identified a family of closely related genes that encodes at least 7 proteins (**Glut-1-Glut-14**, Mol. Wt. 40-80 kDa) and Sodium glucose co-transporter-1 (SGLT-1, 662 amino acids; ~75 kDa). Individual member of this family have identical predicted secondary structures with 12 transmembrane domains. Both N and c-termini are predicted to be cytoplasmic. Most differences in sequence homology exist within the four hydrophilic domains that may play a role in tissue-specific targeting.

Human Glut-1 (GTR1, 492 aa, mol wt ~50-55 kDa, chromosome 1p35-p31.3) belongs to Solute carrier family 2, facilitated glucose transporter member 1 (Slc2A1, Glucose transporter type 1, erythrocyte/brain) or HepG2 glucose transporter. It is a cell membrane; multi-pass membrane protein localizes primarily at the cell surface. Glut-1 has a very broad substrate specificity; can transport a wide range of aldoses including both pentoses and hexoses. Glut-1 mediates glucose transport into red cells, and throughout the blood brain barrier, and supply glucose to most cells.

**Source of Antigen and Antibodies**

<b>Antigen</b>	15-aa peptide from <b>Human Glut-1</b> (gene accession # <a href="#">P11166</a> Designation (GT14-P, control peptide) conjugated to KLH
<b>Location</b>	~N-terminus, Extracellular domain
<b>Ab Host/type</b>	Rabbit, Polyclonal Aff pure IgG (cat # GT14-A) purified over antigen-agarose column
<b>2-ab</b>	<b>Goat Anti-rabbit IgG-HRP</b> cat # 20320 (AP, biotin, FITC conjugates also available)
<b>-ve control</b>	<b># 20009-1, Rabbit (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control</b>

**Recommended Usage**

**Western Blotting** (1-10 ug/ml for affinity pure antibody using ECL technique). (refs 2)

**ELISA:** Control peptide can be used to coat ELISA plates at 1 ug/ml and detected with antibodies (0.5-1 ug/ml for affinity pure).

**Histochemistry & Immunofluorescence:** (see refs 2). We recommend the use of aff pure IgG at 2-20 ug/ml.

**Specificity & Cross-reactivity**

The GT14-P control peptide shows 100% homology in human, rat

(80%), rabbit (86%), pig (80%), and 73% in bovine and mouse. No significant homology is detected with other species. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity (see detailed protocol at: [www.4adi.com\data\abblock.html](http://www.4adi.com\data\abblock.html)).

**General References:** Haspel et al., (1986) J. Biol. Chem. 263, 398-403; Birnbaum, et al., (1986) 83, 5784-5788. 2. Piper et al., (1991) Am. J. Physiol. 260, C570-C580. 3. Harris et al. (1992) Proc. Natl. Acad. Sci. 89, 7556-7560.; see reviews by Baldwin, SA (1993) Biochem. Biophys. Acta 1154, 17-49; Mueckler, M (1994) Eur. J. Biochem. 219, 713-725.

**(2) Citations of ADI's Antibodies for NRAMP2**

Please search for *Gllut1* and consult the publications for additional guidance on antibody concn, and other technique.

[www.4adi.com/serach.php](http://www.4adi.com/serach.php)

\*This product is for In vitro research use only.

**Form & Storage of Antibodies/Peptide Control**

**Antiserum (unpurified)**

100ul solution lyophilized powder  
Supplied in Buffer: 0.05% azide  
**Reconstitute** powder in 100 ul PBS

**Affinity pure IgG**

100 ug/100ul solution lyophilized powder  
Supplied in **Buffer:** PBS+0.1% BSA  
**Reconstitute** powder in PBS at 1mg/ml

**Control/blocking peptide**

100 ug/100 ul solution lyophilized powder  
Supplied in Buffer: PBS pH 7.5,  
**Reconstitute** powder in PBS at 1 mg/ml.

**Storage**

**Short-term:** unopened, undiluted liquid vials at -200C and powder at 4oC or -20oC..

**Long-term:** at -20C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

**Stability:** 6-12 months at -20oC or below.

**Shipping:** 4oC for solutions and room temp for powder

**Related material available from ADI**

Antibodies and Peptides: Glucose transport family (GT 1-13).

GT14-A-P

70501A