

Product Specification Sheet

Hyperpolarization-activated Cyclic Nucleotide-gated channel 2 (HCN2) Antibodies

| | | |
|---------------|---------------------------------------|---------------------|
| Cat # HCN21-S | Rabbit Anti-Human HCN2 antiserum | SIZE: 100 ul |
| Cat # HCN21-A | Rabbit Anti-Human HCN2 IgG (Aff pure) | SIZE: 100 ug |
| Cat # HCN21-P | Human HCN2 Control/blocking peptide | SIZE: 100 ug |

Biological rhythms (beating of the heart, circadian sleep cycles, respiration, and the release of hormones) are necessary to sustain life. Heart beating is the most reliable and rhythmic biological phenomenon. Cardiac pacemaking is produced by the slow diastolic depolarization phase of the action potential. The hyperpolarization-activated cation current (termed I_f , I_h , or I_q) plays a key role in the initiation and modulation of cardiac and neuronal pacemaker depolarization. The generation of cardiac pacemaker potentials relies on a complex interplay between at least four different types of cation channels: T- and L-type Ca^{2+} channels, K^+ channels, and a cation channel termed I_f (synonymous names are I_h and I_q). The I_f channel has been designated as "pacemaker" channel because it reveals unique features that are believed to be a prerequisite for pacemaker activity. Recently, the **hyperpolarization-activated cyclic nucleotide-gated families of ion channel proteins (HCN1-4)** have been identified as the "pacemaker" channel. The amino acid sequences of HCNs predict a structure similar to that of voltage-gated (Kv) channels and cyclic nucleotide-gated (CNG) channels. HCNs proteins (HCN1-4) are characterized by six transmembrane domains (S1-S6), including a including a positively charged voltage-sensing S4 segment and an ion-conducting pore between S5 and S6. In the C terminus the HCNs carry a cyclic nucleotide-binding domain (CNBD), a motif found in several cyclic nucleotide-binding proteins. The core region of HCNs channels (S1 to the C terminus of the CNBD) is highly conserved, whereas the cytoplasmic N and C-termini, vary considerably in their length and share only weak sequence homology.

HCN2, also known as Brain cyclic nucleotide channel gated 2, BCNG2 or HAC-1 (rat 834 aa, human 889-aa, chromosome 19p13.3), is primarily expressed in brain. In mouse, HCN1 is mainly expressed in brain (olfactory bulb and cerebral cortex, hippocampus, thalamus, cerebellum etc) and heart (ventricles and atrium).

Source of Antigen and Antibodies

| | |
|------------------------|--|
| Antigen | 15-aa peptide from human HCN2 (1) ; Designation (#HCN21-P, control/blocking peptide) conjugated to KLH; epitope location ~ N-terminal, Cytoplasmic domain |
| Ab Host/type | Rabbit, Polyclonal unpurified antiserum (#HCN21-S) and IgG, purified over antigen-agarose (Cat # HCN21-A) |
| 2-Ab | Cat # 20320, goat anti-rabbit IgG-HRP (AP, biotin, FITC conjugates also available). |
| -ve control IgG | # 20009-1, Rabbit (non-immune) IgG, purified, suitable for ELISA, Western, IHC as -ve control |

Form & Storage of Antibodies/Peptide Control

Antiserum (unpurified)

100 ul Solution Lyophilized
Supplied 0.1% azide, **Reconstitute** powder in 100 ul PBS

Affinity pure IgG

100 ug/100 ul Solution Lyophilized
Supplied in **Buffer:** PBS + 0.1% BSA
Reconstitute powder in PBS at 1 mg/ml

Control/blocking peptide

100 ug/100 ul Solution Lyophilized
Supplied in **Buffer:** PBS pH 7.5
Reconstitute powder in PBS at 1 mg/ml

Storage

Short-term: unopened, undiluted liquid vials at -20°C and powder at 4°C or -20°C..

Long-term: at -20°C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at -20°C or below.

Shipping: 4°C for solutions and room temp for powder

Recommended Usage

Western Blotting (1:1K-5K for neat serum and 1-10 ug/ml for affinity pure using Chemiluminescence technique). NOT TESTED

ELISA (1:10K-1:100K; using 50-100 ng of control peptide/well).

Histochemistry & Immunofluorescence: NOT TESTED. We recommend the use of affinity pure antibody at 2-20 ug/ml.

Specificity & Cross-reactivity

The HCN21-P peptide is 100% conserved in human, mouse and rat HCN2. No significant sequence homology of HCN21-P is seen with other HCNs or any other protein. Antibody reactivity in various species is not known. The HCN21-P Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity (see detailed protocol at the web site).

General References: (1) Monteggia LM et al (2000) Brain Res. Mol. Brain. Res. 81, 129-139; Ludwig A et al (1999) EMBO J. 18, 2323-2329; Santoro B et al (1997) PNAS 94, 14815-14820; Santoro B et al (1998) Cell 93, 717-729; Waigner BJ et al (2001) Nature 411, 825-810;

**This product is for In vitro research use only.*

HCN21-S-A-P

rev. 201103AC