ALPHA DIAGNOSTIC INTERNATIONAL

Product Specification Sheet

OREXIN-2 Receptor (Hypocretin-2 Receptor) Antibodies

Cat. OX2R22-S	Rabbit Anti-Hhuman Orexin-2 Receptor Antiserum #2	**SIZE:** 100 ul
Cat. OX2R22-A	Rabbit Anti-Human Orexin-2 Receptor IgG #2 (aff pure)	**SIZE**: 100 ug
Cat. OX2R22-P	Human Orexin-2 Receptor Control peptide #2	**SIZE**: 100 ug

Several peptides associated with feeding behavior have been reported recently. Orexins (**Orexin-A** and **Orexin-B**) are a family of hypothalamic neuropeptides selectively expressed in the hypothalamus (1-2). Orexin-A and Orexin-B are derived from the same precursor (Prepro-orexin) by proteolytic cleavage. **Prepro-orexin** is 130 amino acid long peptide with a putative 33-AA secretory sequence, a hydrophobic core followed by residues with small polar side chains. The expression was detected in brain and to a small extent in testis (1-2). These neuropeptides bind and activate two closely related **Orexin receptors**--G-protein coupled receptors (GPCRs) **OX1R** and **OX2R**. Rat and human OX2R are 460 aa, and 444 aa, respectively (1-2)

Source of Antigen and Antibodies

Antigen	19-aa peptide from **human OX2R/hcrtr2 (**gene accession # **O43614, refs 1)**; Designation **(#OX2R22-P, control/blocking peptide**) conjugated to KLH
Epitope Location	~N-terminus Extracellular domain
Ab Host/type	Rabbit, Polyclonal unpurified antiserum (**#OX2R22**-S) and IgG, purified over antigen-agarose (Cat # **OX2R22**-A)
2-Ab	Cat # 20320, goat anti-rabbit IgG-HRP (AP, biotin, FITC conjugates also available).
-ve control	# 20009-1, Rabbit (non-immune) IgG, purified, suitable for ELISA, Western, IHC as –ve control

Form & Storage of Antibodies/Peptide Control

Antiserum (unpurified)
100ul solution lyophilized powder
Supplied in Buffer: 0.05% azide
Reconstitute powder in 100 ul PBS

Affinity pure IgG
100 ug/100ul solution lyophilized powder
Supplied in **Buffer**: PBS+0.1% BSA
Reconstitute powder in PBS at 1mg/ml

Control/blocking peptide
100 ug/100 ul solution lyophilized powder
Supplied in Buffer: PBS pH 7.5,
Reconstitute powder in PBS at 1 mg/ml.

Storage
Short-term: unopened, undiluted liquid vials at -20OC and powder at 4oC or -20oC..

Long-term: at –20C or below in suitable aliquots after reconstitution. Do not freeze and thaw and store working, diluted solutions.

Stability: 6-12 months at –20oC or below.

Shipping: 4oC for solutions and room temp for powder

Recommended Usage

Western Blotting (1:1K-5K for neat serum and 1-10 ug/ml for affinity pure using ECL). (see published refs using this antibody in 2).

ELISA (1:10K-1:100K; using 50-100 ng of control peptide/well).

Histochemistry: We recommend the use of affinity-purified antibody at 2-20 ug/ml (see published refs. 2)

Specificity & Cross-reactivity

Human OX2R22-P peptide is 94% in dog and pig, 84% in rat and 78% in mouse OX2R. No significant homology is seen with OX1R or other G-protein-coupled receptors. Antibody reactivity in all species is not established. Control peptide, because of its low mol. Wt (<3 kDa), is not suitable for Western. It should be used for ELISA or antibody blocking experiments (use 5-10 ug control peptide per 1 ug of aff pure IgG or 1 ul antiserum) to confirm antibody specificity (see detailed protocol at: www.4adi.com\data\abblock.html).

General References: (1) Sakurai, T. et al. (1998) *Cell*, **92**, 573-585; DeLecca L et al (1998) PNAS 95, 322-327

(2) Citations of ADI's Antibodies (see web site for updated list)

Beiras-Fernández A, 2004, J Anatomy 204, 117-122, IHC,
Naslun E, 2002, AJP Gastrointest Liver Physiol 282: G470-G479
Blanco, M., 2001, J. Clin. Endocrinol. Metab. 86: 3444
Blancoa, M, 2001, Regulatory Peptide in press
Beiras-Fernández A, 2004, Journal of Anat. 204, 117
Caillol M 2003 Brain Res. 960, 48-61
Lopez M,1999, Endocrinol. 140, 5991-5994
Blanco M, 2001, J Clin. Endocrinol. Metabol. 86, 1616
Kirchgessner, K., 1999, Neuron 24: 941-951

*This product is for *in vitro* research use only.

Related material available from ADI:

OX2R22-S-A-P 70622A

Alpha Diagnostic Intl Inc., 6203 Woodlake Center Dr, San Antonio, TX 78244, USA; **Email**: service@4adi.com
☎ (800) 786-5777; (210) 561-9515; **Fax**: (210) 561-9544; **Web Site**: www.4adi.com