



Product Data Sheet

Cat # RP-1035

Recombinant Human MIG (CXCL9)

Size: 5 ug

Chemokine (C-X-C motif) ligand 9 (CXCL9) is a small cytokine belonging to the CXC chemokine family that is also known as Monokine induced by gamma interferon (MIG). CXCL9 is a T-cell chemo attractant, which is induced by IFN- γ . It is closely related to two other CXC chemokines called CXCL10 and CXCL11, whose genes are located near the gene for CXCL9 on human chromosome 4. CXCL9, CXCL10 and CXCL11 all elicit their chemotactic functions by interacting with the chemokine receptor CXCR3.

Source: *Escherichia Coli*. MIG (monokine induced by gamma-interferon) Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 103 amino acids and having a molecular mass of 11700 Dalton. The MIG is purified by proprietary chromatographic techniques. The CXCL9 was lyophilized from a concentrated (1mg/ml) solution in water containing no additives.

Applications and Suggested Dilutions: Greater than 95.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE. It is recommended to reconstitute the lyophilized MIG in sterile 18M Ω -cm H₂O not less than 100 μ g/ml, which can then be further diluted to other aqueous solutions. Users must optimize the appropriate concentration and conditions for each assay.

Storage and Stability: Lyophilized MIG although stable at room temperature for 3 weeks, should be stored desiccated below -18 $^{\circ}$ C. Upon reconstitution CXCL9 should be stored at 4 $^{\circ}$ C between 2-7 days and for future use below -18 $^{\circ}$ C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). If supplied in powder then reconstitute it in 100 ul water for 1 mg/ml stock and store in liquid at 4 $^{\circ}$ C for ~1 week or aliquots in suitable size and store at -20 $^{\circ}$ C for long term storage. **Please prevent freeze-thaw cycles.**

Biological Activity: ED₅₀ range=10.0-100.0 ng/mL, determined by the dose dependent chemotaxis with human lymphocytes cultured in the presence of IL-

Usage:

This item is for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals

RP-1035 120419V