



Product Data Sheet

□ Cat # RP-386

Recombinant Human Paraoxonase-2

Size: □ 10 ug

Paraoxonase 2 (PON2) is a member of a multigene family whose genes share 65% identity at the amino acid level, and is expressed in a variety of tissues, including the pancreas. PON2 overexpression has been shown to lower the intracellular oxidative state and reduce the cells ability to oxidize LDL. PON2 is therefore implicated in the modulation of oxidative stress.

Source: *Escherichia Coli*. Paraoxonase-2 Human Recombinant is expressed in E. coli having a molecular weight of 43.5 kDa and fused to an amino terminal hexahistidine tag. The PON2 purified by proprietary chromatographic techniques. 0.1 mg/ml PON2 is supplied in PBS and 50% glycerol.

Applications and Suggested Dilutions: Arylesterase 2 can be used directly as a positive control in Western blotting, ELISA, immunoprecipitation and other immunological experiments. The biological activity of this product has not yet been tested. Greater than 95% as determined by SDS-PAGE. Single band on Western Blot. Users must optimize the appropriate concentration and conditions for each assay.

Storage and Stability: Store at 4°C if entire vial will be used within 1-2 weeks. Store, frozen at -20°C for longer periods of time. **Avoid multiple freeze-thaw cycles.** If supplied in powder then reconstitute it in 100 ul water for 1 mg/ml stock and store in liquid at 4oC for ~1 week or aliquots in suitable size and store at -20oC for long term storage.

Usage: This item is for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

RP-386

110207V